



DIACOMIT® (stiripentol)

Access and Coverage Guide

Find out how you and your patients/caregivers can **access the unique services only available with brand-name DIACOMIT** through our exclusive specialty pharmacy, PANTHERx Rare.

INDICATION

DIACOMIT (stiripentol) is indicated for the treatment of seizures associated with Dravet syndrome (DS) in patients taking clobazam who are 6 months of age and older and weighing 7 kg or more. There are no clinical data to support the use of DIACOMIT as monotherapy in Dravet syndrome.

IMPORTANT SAFETY INFORMATION

CONTRAINDICATIONS None

DIACOMIT® (stiripentol) Is FIRST and PROVEN

When you choose DIACOMIT, you're offering your patients with Dravet syndrome:

The FIRST Dravet-specific antiseizure medication:



- With nearly **three decades** of real-world use in over 55 countries¹⁻⁵
- FDA approved for children **as young as 6 months***¹

** Weighing 15 lb or more and taking clobazam*

A PROVEN, effective treatment:



- Significantly **reducing seizures** associated with Dravet syndrome^{1,6}
- Helping patients achieve **seizure freedom**^{1,6}
- With 69,000 patient-years of experience⁵

Quality and Consistency You Can Count On

For many patients with Dravet, it's hard to find a treatment that works. That's why consistency is key.

When you prescribe DIACOMIT, your patients/caregivers get the same quality and unique services with every refill.

IMPORTANT SAFETY INFORMATION

WARNINGS & PRECAUTIONS

Somnolence

DIACOMIT can cause somnolence. Monitor patients for somnolence, particularly when DIACOMIT is used concomitantly with other CNS depressants or clobazam, which is also known to cause somnolence.

Decreased Appetite and Decreased Weight

DIACOMIT can cause decreases in appetite and weight. The growth and weight of pediatric patients treated with DIACOMIT should be carefully monitored.

Neutropenia and Thrombocytopenia

DIACOMIT can cause significant declines in neutrophil and platelet counts. Hematologic testing should be obtained prior to starting treatment with DIACOMIT and then every 6 months.

Biocodex Partners With PANTHERx Rare Pharmacy

DIACOMIT is available **exclusively through PANTHERx Rare Pharmacy**, a national specialty pharmacy that manages the prescription process and provides you and your patients/caregivers with concierge customer service.



BIOCODEX
by your side

**NO
REMS**

With **no Risk Evaluation and Mitigation Strategy (REMS)** requirements for DIACOMIT, PANTHERx can deliver treatment directly to your patient's door in **as little as 24 hours**.

PANTHERx Services and Programs To Support Your Patients

With **brand-name DIACOMIT**, you and your patients/caregivers will maintain access to personalized, exclusive services through our specialty pharmacy partner.



For healthcare professionals, the Biocodex By Your Side Support Team at PANTHERx will:

- Help you **navigate prior authorization/reauthorization** and appeals processes
- Conduct **full benefit investigations** and confirm patient eligibility for savings
- **Proactively contact you** for additional refills once the final prescription has been filled



Exclusive concierge services for your patients and their caregivers:

- A **dedicated team** that understands Dravet syndrome and treatment with DIACOMIT
- Pharmacists **on call 24/7** and translation services in 100+ languages
- A Rapid Refill program to help patients **conveniently schedule refills online** or via mobile (opt-in required)
- **Shipment tracking** and **vacation planning**, so your patients/caregivers know when to expect their delivery and do not miss a dose
- **Dosing supplies** by request (10 mL and 35 mL oral syringes as well as 100 mL dosing cups)

WARNINGS & PRECAUTIONS

Withdrawal Symptoms As with most antiepileptic drugs (AEDs), DIACOMIT should be gradually withdrawn to minimize the risk of increased seizure frequency and status epilepticus.

Biocodex Is Putting Patients First

No matter the circumstances, Biocodex has your patients with Dravet covered. There are four patient access programs in place to help ensure that your patients can easily access treatment, regardless of their financial status or insurance coverage.

Our Patient Savings and Support Programs

The Biocodex By Your Side Support Team at PANTHERx evaluates all patients' eligibility for the following programs.



Copay Program

- **\$0 copay** per prescription
- For eligible patients with commercial insurance
- Not available for government plans such as Medicare or Medicaid



Quick Start Program

- Up to 60-day fulfillment of DIACOMIT at **no cost** to the patient
- For newly eligible patients experiencing delays in gaining access to treatment



Bridge Program

- Up to 60-day fulfillment of DIACOMIT at **no cost** to the patient
- For patients on DIACOMIT with changes or lapses in their insurance coverage



Long-Term Patient Assistance Program

- Prescription of DIACOMIT at **no cost** to the patient
- For eligible patients who are uninsured or underinsured

Eligible patients with commercial insurance may pay as little as \$0 for DIACOMIT.

WARNINGS & PRECAUTIONS

Risks in Patients with Phenylketonuria (PKU)

DIACOMIT for oral suspension contains phenylalanine, which can be harmful to patients with PKU. Before prescribing DIACOMIT for oral suspension to a patient with PKU, consider the total daily intake of phenylalanine from all sources, including DIACOMIT for oral suspension. DIACOMIT capsules do not contain phenylalanine.

Suicidal Behavior and Ideation

AEDs, including DIACOMIT, increase the risk of suicidal thoughts or behavior. Patients treated with any AED for any indication should be monitored for the emergence or worsening of depression, suicidal thoughts or behavior, and/or any unusual changes in mood or behavior.

Two Ways To Get Started

1. Send a Completed Patient Enrollment Form via Fax to PANTHERx at 833-871-4137

OR

2. Send an e-prescription to PANTHERx via an Electronic Health Record (EHR)*

1120 Stevenson Mill Road, Suite 400,
Coraopolis, PA 15108

**Remember to add DIACOMIT as a "Favorite Medication" within your EHR to simplify the ordering process.*

To avoid medication delays please:

- ✓ Remember to include:
 - Diagnosis code(s) and patient's weight
 - Formulation, with clear dosing/titration instructions, and quantity (eg, number of capsules)
 - A copy of your patient's prescription and medical benefit cards (front and back), if utilizing the enrollment form
 - List of your patient's current, discontinued or failed medications, including clobazam use and dosage
 - Your patient's medical records
 - An office contact name and phone number
- ✓ Request or sign that **brand-name DIACOMIT** must be dispensed and no generic substitution is allowed (see pages 6-7 for state-specific prescription requirements)



DIACOMIT is only available through PANTHERx Rare Pharmacy

With **brand-name DIACOMIT**, you and your patients/caregivers will maintain access to personalized, exclusive services through our specialty pharmacy partner.

ADVERSE REACTIONS

The most common adverse reactions that occurred in at least 10% of DIACOMIT-treated patients and more frequently than on placebo were somnolence, decreased appetite, agitation, ataxia, decreased weight, hypotonia, nausea, tremor, dysarthria, and insomnia.

PREGNANCY

There are no adequate data on the developmental risks associated with the use of DIACOMIT in pregnant women. Based on animal data, DIACOMIT may cause fetal harm. There is a pregnancy exposure registry that monitors pregnancy outcomes in women exposed to AEDs, such as DIACOMIT, during pregnancy. Physicians are advised to recommend that pregnant patients taking DIACOMIT enroll in the North American Antiepileptic Drug (NAAED) Pregnancy Registry (information at www.aedpregnancyregistry.org). This can be done by calling the toll-free number 1-888-233-2334 and must be done by patients themselves or their caregiver.

State “Dispense As Written” Requirements*7

If you and your patients/caregivers decide to treat with DIACOMIT® (stiripentol), the chart below can help you determine special instructions required by your state or territory to help ensure that your patient receives brand-name DIACOMIT as intended.

State	Requirements To Prevent Substitution
Alabama	Prescriber must expressly indicate in some manner
Alaska	Must write in own handwriting, in addition to signature, “Brand Necessary”
Arizona	Prescriber must expressly indicate that substitution is not allowed
Arkansas	Must write in own handwriting, in addition to signature, “Brand Necessary”
California	Prescriber may indicate orally or in own handwriting “Do Not Substitute” or similar words; allows use of preprinted “Do Not Substitute” box if the prescriber personally initials the box
Colorado	Prescriber must handwrite “Dispense as Written” or hand initial a preprinted box labeled “Dispense as Written”; may also be done electronically
Connecticut	Prescriber must write in own handwriting “DAW” or “Dispense as Written” and “Medically Necessary”
Delaware	Prescriber must write in own handwriting “DAW,” “Dispense as Written,” “Brand necessary,” or “Brand medically necessary”
District of Columbia	Prescriber must expressly indicate in some manner
Florida	Prescriber must expressly indicate in some manner
Georgia	Prescriber’s signature shall validate the prescription and, unless the prescriber handwrites “Brand Necessary” or “Brand Medically Necessary,” shall designate approval of drug substitution by the pharmacist
Guam	A licensed practitioner shall prohibit drug product selection by handwriting the words “No Substitution” or the diminutive “No Sub” on the face of the prescription
Hawaii	Prescriber must indicate “brand necessary” or “brand medically necessary” in own handwriting or product selection is allowed; refer to the Department of Health, Food, and Drug Branch
Idaho	If a prescriber orders by any means that a brand-name drug must be dispensed, then no drug selection is permitted
Illinois	Prescriber must indicate “may not substitute” by marking a designated box; see Section 225 ILCS 85/25
Indiana	Prescriber’s signature required on appropriate line of two-line prescription
Iowa	Prescriber must expressly indicate that substitution is not allowed
Kansas	Prescriber’s signature required on appropriate line of two-line prescription; prescriber must expressly indicate that substitution is not allowed
Kentucky	Prescriber must expressly indicate in some manner; for Medicaid patients, “Brand Medically Necessary” to be handwritten on the face of the prescription by the prescriber, or product selection is allowed; May indicate in manner of his or her choice on the prescription “Do Not Substitute,” except that the indication shall not be preprinted on a prescription
Louisiana	Box must be checked
Maine	Prescriber must expressly indicate in some manner; box must be checked
Maryland	Prescriber must expressly indicate that substitution is not allowed; for Medicaid patients, “Brand Medically Necessary” to be handwritten on the face of the prescription by the prescriber, or product selection is allowed
Massachusetts	Prescriber must indicate “No Substitution”
Michigan	Prescriber must write in own handwriting “DAW” or “Dispense as Written”; prescriber expressly indicates that prescription is to be dispensed as communicated for prescriptions other than those written
Minnesota	Prescriber must write in own handwriting “DAW” or “Dispense as Written” unless the prescription is transmitted electronically in accordance with Code of Federal Regulations, Title 42, Section 423
Mississippi	Prescriber’s signature required on appropriate line of two-line prescription
Missouri	If a prescriber orders by any means that a brand-name drug must be dispensed, then no selection is permitted
Montana	The prescriber must indicate “Brand name medically necessary” on the prescription if it is medically necessary that an equivalent drug or biologic product not be selected

State “Dispense As Written” Requirements (cont.)*7

State	Requirements To Prevent Substitution
Nebraska	Prescriber must expressly indicate in some manner
Nevada	Prescriber must write in own handwriting “Dispense as Written”
New Hampshire	Prescriber must handwrite “medically necessary” on each paper prescription, or uses electronic indications when transmitted electronically, or gives instructions when transmitted orally that the brand-name drug product is medically necessary
New Jersey	Prescriber’s signature required on appropriate line of two-line prescription
New Mexico	A licensed practitioner shall prohibit drug product selection by handwriting the words “No Substitution” or the diminutive “No Sub” on the face of the prescription
New York	Prescriber must indicate “Dispense as Written” in the designated box, or positively indicate brand for electronic prescriptions; for Medicaid patients, “Brand Medically Necessary” to be handwritten on the face of the prescription by the prescriber, or product selection is allowed
North Carolina	Prescriber’s signature required on appropriate line of two-line prescription; prescriber must expressly indicate in some manner
North Dakota	Prescriber must write in own handwriting in addition to signature “Brand Medically Necessary”
Ohio	In the case of written or electronic prescription, including computer-generated prescription, the prescriber handwrites or actively causes to display on the prescription “dispense as written,” “DAW,” “do not substitute,” “brand medically necessary,” or any other statement or numerical code that indicates the prescriber’s intent to prevent substitution; such designation shall not be preprinted or stamped on the prescription, but a reminder to the prescriber of the designation procedure may be preprinted or displayed on the form or electronic system used to issue the prescription
Oklahoma	O.S. (1961) states that it is unlawful for a pharmacist to substitute without the authority of the prescriber or purchaser
Oregon	Refer to ORS 689.515 and OAR-855-041-1105
Pennsylvania	Prescriber’s signature shall validate the prescription and, unless the prescriber handwrites “Brand Necessary” or “Brand Medically Necessary,” shall designate approval of drug substitution by the pharmacist
Puerto Rico	Prescriber must write on the face of the prescription in own handwriting the phrase “Do not interchange”
Rhode Island	Prescriber’s signature shall validate the prescription and, unless the prescriber indicates “Brand Necessary” or “Brand Medically Necessary,” shall designate approval of drug substitution by the pharmacist; patient may request, in writing, that the brand name be dispensed
South Carolina	Prescriber’s signature required on appropriate line of two-line prescription
South Dakota	Prescriber must write in own handwriting in addition to signature “Brand Necessary”
Tennessee	The prescriber shall, in the prescriber’s own handwriting, include on the prescription the following language (but not limited to): (1) “brand name medically necessary,” “dispense as written,” “medically necessary,” “brand name,” “no generic,” or (2) any abbreviation of the language in (1), or (3) any other prescriber handwritten notation, such as circling a preprinted “dispense as written” on the prescription order, that clearly conveys the intent that a brand name is necessary for the patient
Texas	Prescriber must indicate “brand necessary” or “brand medically necessary” in own handwriting or product selection is allowed
Utah	Prescriber must expressly indicate in some manner; allows use of preprinted “Do Not Substitute” checkbox
Vermont	Prescriber must write “brand necessary,” “no substitution,” “dispense as written,” or “DAW” in own handwriting (see 18 V.S.A. §4606 Brand Certification)
Virginia	“Brand Medically Necessary” to be handwritten on the face of the prescription by the prescriber for Medicaid patients, or product selection is allowed; for all non-Medicaid patients, phrase must be included but not required to be handwritten
Washington	Prescriber’s signature required on appropriate line of two-line prescription
West Virginia	Prescriber must write in own handwriting “Brand Medically Necessary”
Wisconsin	Prescriber must expressly indicate in some manner
Wyoming	Prescriber must expressly indicate that substitution is not allowed

*State and territory regulations are subject to change. Please consult your state or territory’s Board of Pharmacy and Medicaid office to verify prescribing requirements.

Reprinted with permission from the National Association of Boards of Pharmacy® (NABP®). Last updated 2025.

DIACOMIT: Quality and Consistency You Can Count On



Biocodex partners with PANTHERx Rare Pharmacy

They provide exclusive concierge services to streamline the access process.



DIACOMIT can be delivered in as little as 24 hours

- No REMS required
- No known contraindications*

*Clinical discretion is advised when adjusting antiseizure medication regimens.



Eligible patients with commercial insurance may pay as little as \$0

There are additional savings and support programs to help patients access treatment.



Choose brand-name DIACOMIT

With brand-name DIACOMIT, you and your patients/caregivers will maintain access to personalized, exclusive services through PANTHERx Rare Pharmacy.

Contact the Biocodex By Your Side Support Team at PANTHERx



833-248-0467
Monday – Friday, 8 AM – 8 PM E.T.



1120 Stevenson Mill Road, Suite 400
Coraopolis, PA 15108



For helpful resources
and information
about DIACOMIT visit
[DIACOMIT.com/hcp](https://diacomit.com/hcp)

IMPORTANT SAFETY INFORMATION

To report suspected adverse reactions, contact Biocodex at 1-866-330-3050 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Please see full Prescribing Information for DIACOMIT® enclosed or at [DIACOMIT.com](https://diacomit.com).

References: 1. DIACOMIT® [prescribing information]. Beauvais, France: Biocodex, Inc.; July 2022. 2. Chiron C, Marchand MC, Tran A, et al; for the STICLO study group. Stiripentol in severe myoclonic epilepsy in infancy: a randomised placebo-controlled syndrome dedicated trial. *Lancet*. 2000;356(9242):1638-1642. 3. Farwell JR, Anderson GD, Kerr BM, Tor JA, Levy RH. Stiripentol in atypical absence seizures in children: an open trial. *Epilepsia*. 1993 Mar-Apr; 34(2): 305-11. 4. Bebin M, Bleck TP. New anticonvulsant drugs. Focus on flunarizine, fosphenytoin, midazolam and stiripentol. *Drugs*. 1994 Aug; 48(2): 153-71. 5. DIACOMIT Periodic Benefit-Risk Evaluation Reports (PBRERs). Beauvais, France: Biocodex, Inc.; January 2022. 6. U.S. Food and Drug Administration. CDER Clinical Review. August 2018. https://www.accessdata.fda.gov/drugsatfda_docs/nda/2018/206709Orig1s000,207223Orig1s000MedR.pdf. Accessed May 12, 2020. 7. National Association of Boards of Pharmacy. 2025 Survey of Pharmacy Law. Mount Prospect, IL; 2025:110-114.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use DIACOMIT safely and effectively. See full prescribing information for DIACOMIT.

DIACOMIT (stiripentol) capsules, for oral use

DIACOMIT (stiripentol) for oral suspension

Initial U.S. Approval: 2018

-----RECENT MAJOR CHANGES-----

Indications and Usage (1) 7/2022

Dosage and Administration (2.2) 7/2022

-----INDICATIONS AND USAGE-----

DIACOMIT is indicated for the treatment of seizures associated with Dravet syndrome in patients taking clobazam who are 6 months of age and older and weighing 7 kg or more. There are no clinical data to support the use of DIACOMIT as monotherapy in Dravet syndrome. (1)

-----DOSAGE AND ADMINISTRATION-----

- The dosage of DIACOMIT is 50 mg/kg/day, administered by mouth in 2 or 3 divided doses, depending on age and weight. (2.2)
- Capsules must be swallowed whole with a glass of water during a meal. Capsules should not be broken or opened. (2.3)
- Powder for suspension should be mixed in a glass of water and should be taken immediately after mixing during a meal. (2.3)
- Reduce dose or discontinue dose gradually. (2.5)

-----DOSAGE FORMS AND STRENGTHS-----

- Capsule: 250 mg or 500 mg (3)
- For Oral Suspension: 250 mg or 500 mg (3)

-----CONTRAINDICATIONS-----

None (4)

-----WARNINGS AND PRECAUTIONS-----

- *Somnolence:* Monitor for somnolence, particularly when DIACOMIT is used concomitantly with other CNS depressants; If somnolence occurs

during co-administration with clobazam, consider an initial reduction of clobazam by 25%. (5.1)

- *Decreased Appetite and Decreased Weight:* the weight of patients and the growth rate of pediatric patients should be carefully monitored. (5.2)
- *Neutropenia and Thrombocytopenia:* Blood counts should be obtained prior to starting treatment with DIACOMIT and then every 6 months. (5.3)
- *Withdrawal:* DIACOMIT should be gradually withdrawn to minimize the risk of increased seizure frequency and status epilepticus. (5.4)
- *Risks in Patients with Phenylketonuria (PKU):* DIACOMIT for oral suspension contains phenylalanine; consider total daily intake before prescribing to patients with PKU. (5.5)
- *Suicidal Behavior and Ideation:* Monitor for suicidal thoughts or behaviors. (5.6)

-----ADVERSE REACTIONS-----

Adverse reactions that occurred in at least 10% of DIACOMIT-treated patients and more frequently than on placebo were somnolence, decreased appetite, agitation, ataxia, weight decreased, hypotonia, nausea, tremor, dysarthria, and insomnia. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact BIOCDEX at 1-866-330-3050 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

-----DRUG INTERACTIONS-----

- DIACOMIT increases the plasma concentration of clobazam and its metabolite through metabolic inhibition of CYP3A4 and CYP2C19. Consider dose reduction of clobazam in case of adverse reactions. (7.1)
- Substrates of CYP2C8, CYP2C19, P-gp and BCRP may require a dose reduction. (7.1)
- Substrates of CYP1A2, CYP2B6 and CYP3A4 may require a dose adjustment. (7.1)
- Strong inducers of CYP1A2, CYP3A4 or CYP2C19: Consider dose increase of DIACOMIT. (7.2)

-----USE IN SPECIFIC POPULATIONS-----

- *Pregnancy:* Based on animal data, may cause fetal harm. (8.1)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.

Revised: 7/2022

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Laboratory Tests Prior to First Dose of DIACOMIT
- 2.2 Dosing Information
- 2.3 Important Administration Instructions
- 2.4 Missed Dose
- 2.5 Gradual Withdrawal

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Somnolence
- 5.2 Decreased Appetite and Decreased Weight
- 5.3 Neutropenia and Thrombocytopenia
- 5.4 Withdrawal Symptoms
- 5.5 Risks in Patients with Phenylketonuria
- 5.6 Suicidal Behavior and Ideation

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience

7 DRUG INTERACTIONS

- 7.1 Effect of DIACOMIT on Other Drugs
- 7.2 Effect of Other Drugs on DIACOMIT
- 7.3 CNS Depressants and Alcohol

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment
- 8.7 Hepatic Impairment

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

- 16.1 How Supplied
- 16.2 Storage and Handling

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

DIACOMIT is indicated for the treatment of seizures associated with Dravet syndrome (DS) in patients taking clobazam who are 6 months of age and older and weighing 7 kg or more. There are no clinical data to support the use of DIACOMIT as monotherapy in Dravet syndrome.

2 DOSAGE AND ADMINISTRATION

2.1 Laboratory Tests Prior to First Dose of DIACOMIT

Hematologic testing should be obtained prior to starting treatment with DIACOMIT [see *Warnings and Precautions* (5.3)].

2.2 Dosing Information

The recommended oral dosage of DIACOMIT is 50 mg/kg/day, administered in 2 or 3 divided doses (i.e., 16.67 mg/kg three times daily or 25 mg/kg twice daily), depending on the patient's age and body weight as shown in Table 1. If the exact dosage is not achievable given the available strengths, round to the nearest possible dosage, which is usually within 50 mg to 150 mg of the recommended 50 mg/kg/day. A combination of the two DIACOMIT strengths can be used to achieve this dosage. The maximum recommended total dosage is 3,000 mg/day.

Table 1. Recommended Dosage for Patients 6 Months of Age and Older Weighing 7 kg or More with Dravet Syndrome

Age of Patient	Body Weight	Dosing Regimen (administered by mouth in equally divided doses)	Total Daily Dose
6 months to less than 1 year	7 kg and above	25 mg/kg twice daily ^{a,b}	50 mg/kg/day
1 year and above	7 kg to less than 10 kg	25 mg/kg twice daily ^b	50 mg/kg/day
	10 kg and above	25 mg/kg twice daily or 16.67 mg/kg three times daily	50 mg/kg/day Maximum daily dose is 3000 mg

^a Dosing frequency should not exceed twice daily to limit free water administration.

^b Dosing frequency should not exceed twice daily to avoid overexposures.

2.3 Important Administration Instructions

DIACOMIT Capsules

DIACOMIT capsules must be swallowed whole with a glass of water during a meal. Capsules should not be broken or opened.

DIACOMIT for Oral Suspension

DIACOMIT should be mixed in a glass of water (100 mL) and should be taken immediately after mixing during a meal. To be sure there is no medicine left in the glass, add a small amount of water (25 mL) to the drinking cup and drink all of the mixture [see *Instructions for Use*].

2.4 Missed Dose

A missed dose should be taken as soon as possible. If it is almost time for the next dose, the missed dose should not be taken. Instead, the next scheduled dose should be taken. Doses should not be doubled.

2.5 Gradual Withdrawal

As is advisable for most antiepileptic drugs, if DIACOMIT treatment is discontinued, the drug should be withdrawn gradually to minimize the risk of increased seizure frequency and status epilepticus [see *Warnings and Precautions* (5.4)].

In situations where rapid withdrawal of DIACOMIT is medically required, appropriate monitoring is recommended.

3 DOSAGE FORMS AND STRENGTHS

Capsules

- 250 mg: size 2, pink, and imprinted with “Diacomit” and “250mg”
- 500 mg: size 0, white, and imprinted with “Diacomit” and “500mg”

For Oral Suspension

- Pale pink fruit flavored powder packaged in packets. Each packet contains either 250 mg or 500 mg of stiripentol

4 CONTRAINDICATIONS

None.

5 WARNINGS AND PRECAUTIONS

5.1 Somnolence

DIACOMIT can cause somnolence. In controlled studies in patients with Dravet syndrome, the incidence of somnolence was 67% in DIACOMIT-treated patients, compared to 23% in patients on placebo. All patients in both groups were on concomitant clobazam, which is also known to cause somnolence. Co-administration of DIACOMIT with clobazam results in increased levels of clobazam and its active metabolite [see *Drug Interactions* (7.1)]. Other central nervous system CNS depressants, including alcohol, could potentiate the somnolence effect of DIACOMIT.

Prescribers should monitor patients for somnolence. If somnolence occurs during co-administration with clobazam, consider an initial reduction of clobazam by 25%. If somnolence persists, further clobazam reduction by an additional 25% should be considered, as should adjustment of the dosage of other concomitant anticonvulsant drugs with sedating properties. Prescribers should caution patients against engaging in hazardous activities requiring mental alertness, such as operating dangerous machinery or motor vehicles, until the effect of DIACOMIT on mental alertness is known.

5.2 Decreased Appetite and Decreased Weight

DIACOMIT can cause decreases in appetite and weight. In controlled studies in patients with Dravet syndrome, the incidence of decreased appetite was 46% in DIACOMIT-treated patients, compared to 10% in patients on placebo. The incidence of decreased weight was 27% in DIACOMIT-treated patients, compared to 6% in patients on placebo. Nausea and vomiting also occurred more frequently in DIACOMIT-treated patients [*see Adverse Reactions (6.1)*]. Given the frequency of these adverse reactions, the growth of pediatric patients treated with DIACOMIT should be carefully monitored. In some cases, decreasing the dose of concomitant valproate by 30% per week can reduce the decrease in appetite and weight.

5.3 Neutropenia and Thrombocytopenia

DIACOMIT can cause a significant decline in neutrophil count. In controlled studies in patients with Dravet syndrome, there were 31 patients treated with DIACOMIT who had both a baseline and end-of-study neutrophil count obtained. A decrease in neutrophil count from normal at baseline to less than 1500 cells/mm³ during the trial was observed in 13% of these DIACOMIT-treated patients, but not in any placebo-treated patients.

DIACOMIT can cause a significant decline in platelet count. In controlled studies in patients with Dravet syndrome, there were 31 patients treated with DIACOMIT who had both a baseline and end-of-study platelet count. A decrease in platelet count from normal at baseline to less than 150,000/μL during the trial was observed in 13% of these DIACOMIT-treated patients, but not in any placebo-treated patients.

Hematologic testing should be obtained prior to starting treatment with DIACOMIT, and then every 6 months.

5.4 Withdrawal Symptoms

As with most antiepileptic drugs, DIACOMIT should generally be withdrawn gradually to minimize the risk of increased seizure frequency and status epilepticus.

In situations where rapid withdrawal of DIACOMIT is required (e.g., in the setting of a serious adverse reaction), appropriate monitoring is recommended.

5.5 Risks in Patients with Phenylketonuria

Phenylalanine can be harmful to patients with phenylketonuria (PKU). DIACOMIT for oral suspension contains phenylalanine, a component of aspartame. Each 250 mg packet contains 1.40 mg phenylalanine; each 500 mg packet contains 2.80 mg phenylalanine. Before prescribing DIACOMIT for oral suspension to a patient with PKU, consider the combined daily amount of phenylalanine from all sources, including DIACOMIT for oral suspension.

DIACOMIT capsules do not contain phenylalanine.

5.6 Suicidal Behavior and Ideation

AEDs, including DIACOMIT, increase the risk of suicidal thoughts or behavior in patients taking these drugs for any indication. Patients treated with any AED for any indication should be monitored for the emergence or worsening of depression, suicidal thoughts or behavior, and/or any unusual changes in mood or behavior.

Pooled analyses of 199 placebo-controlled clinical trials (mono- and adjunctive therapy) of 11 different AEDs showed that patients randomized to one of the AEDs had approximately twice the risk (adjusted relative risk 1.8, 95% confidence interval [CI]:1.2, 2.7) of suicidal thinking or behavior compared to patients randomized to placebo. In these trials, which had a median treatment duration of 12 weeks, the estimated incidence rate of suicidal behavior or ideation among 27,863 AED treated patients was 0.43%, compared to 0.24% among 16,029 placebo treated patients, representing an increase of approximately one case of suicidal thinking or behavior for every 530 patients treated. There were four suicides in drug treated patients in the trials and none in placebo treated patients, but the number is too small to allow any conclusion about drug effect on suicide.

The increased risk of suicidal thoughts or behavior with AEDs was observed as early as one week after starting drug treatment with AEDs and persisted for the duration of treatment assessed. Because most trials included in the analysis did not extend beyond 24 weeks, the risk of suicidal thoughts or behavior beyond 24 weeks could not be assessed.

The risk of suicidal thoughts or behavior was generally consistent among drugs in the data analyzed. The finding of increased risk with AEDs of varying mechanisms of action and across a range of indications suggests that the risk applies to all AEDs used for any indication. The risk did not vary substantially by age (5-100 years) in the clinical trials analyzed. Table 2 shows absolute and relative risk by indication for all evaluated AEDs.

Table 2. Risk by Indication for Antiepileptic Drugs in the Pooled Analysis

Indication	Placebo Patients with Events per 1000 Patients	Drug Patients with Events per 1000 Patients	Relative Risk: Incidence of Drug Events in Drug Patients/Incidence in Placebo Patients	Risk Difference: Additional Drug Patients with Events per 1000 Patients
Epilepsy	1.0	3.4	3.5	2.4
Psychiatric	5.7	8.5	1.5	2.9
Other	1.0	1.8	1.9	0.9
Total	2.4	4.3	1.8	1.9

The relative risk for suicidal thoughts or behavior was higher in clinical trials for epilepsy than in clinical trials for psychiatric or other conditions, but the absolute risk differences were similar for the epilepsy and psychiatric indications.

Anyone considering prescribing DIACOMIT or any other AED must balance the risk of suicidal thoughts or behavior with the risk of untreated illness. Epilepsy and many other illnesses for which AEDs are prescribed are themselves associated with morbidity and mortality and an increased risk of suicidal thoughts and behavior. Should suicidal thoughts and behavior emerge during treatment, the prescriber needs to consider whether the emergence of these symptoms in any given patient may be related to the illness being treated.

Patients, their caregivers, and families should be informed that AEDs increase the risk of suicidal thoughts and behavior and should be advised of the need to be alert for the emergence or worsening of the signs and symptoms of depression, any unusual changes in mood or behavior, or the emergence of suicidal thoughts, behavior, or thoughts about self-harm. Behaviors of concern should be reported immediately to healthcare providers.

6 ADVERSE REACTIONS

The following serious or otherwise clinically significant adverse reactions are described elsewhere in the labeling:

- Somnolence [*see Warnings and Precautions (5.1)*]
- Decreased Appetite and Decreased Weight [*see Warnings and Precautions (5.2)*]
- Neutropenia and Thrombocytopenia [*see Warnings and Precautions (5.3)*]
- Withdrawal Symptoms [*see Warnings and Precautions (5.4)*]
- Risks in Patients with Phenylketonuria [*see Warnings and Precautions (5.5)*]
- Suicidal Behavior and Ideation [*see Warnings and Precautions (5.6)*]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in clinical trials of another drug, and may not reflect the rates observed in practice.

During its development for the treatment of seizures associated with Dravet syndrome, DIACOMIT was administered to 55 healthy male volunteers and 438 patients with Dravet syndrome, including 310 patients treated for 12 months or more. The conditions and duration of exposure varied greatly, and included single- and multiple-dose clinical pharmacology studies in healthy male volunteers, 2 randomized, double-blind, placebo-controlled, 12-week studies in patients with Dravet syndrome (Study 1 and Study 2), and open-label long-term studies.

In Study 1 and Study 2, 33 patients received DIACOMIT and 31 patients received placebo for a treatment duration of 8 weeks [*see Clinical Studies (14)*]. Adverse reactions from these trials are presented below. Approximately 53% of patients were female and the mean age was 9.2 years. All patients were taking clobazam and valproate.

There were 2 patients in whom adverse reactions led to discontinuation of DIACOMIT treatment: one patient had an adverse reaction of status epilepticus; the second patient had drowsiness, balance impaired and sialorrhea.

The most common adverse reactions, occurring in at least 10% of DIACOMIT-treated patients and more frequently than on placebo, included somnolence (67%), decreased appetite (45%), agitation (27%), ataxia (27%), weight decreased (27%), hypotonia (24%), nausea (15%), tremor (15%), dysarthria (12%), and insomnia (12%).

Table 3 lists the adverse reactions that occurred in 5% or more of DIACOMIT-treated patients and at a rate greater than in patients on placebo in the 2 randomized, double-blind, placebo-controlled, clinical trials in patients with Dravet syndrome (Study 1 and Study 2).

Table 3. Adverse Reactions in 5% or More of DIACOMIT-Treated Patients and More Frequently than on Placebo in Patients with Dravet Syndrome (Study 1 and Study 2)

Adverse Reactions	Study 1 and 2 – Pooled Total	
	DIACOMIT (50mg/kg/day) N=33 %	Placebo N=31 %
Gastrointestinal disorders		
Nausea	15	3
Vomiting	9	0
Salivary hypersecretion	6	0
General disorders and administration site conditions		
Fatigue	9	3
Pyrexia	6	3
Infections and infestations		
Bronchitis	6	0
Nasopharyngitis	6	0
Investigations		
Weight decreased	27	6
Weight increased	6	3
Metabolism and nutrition disorders		
Decreased appetite	46	10
Nervous system disorders		
Somnolence	67	23
Ataxia	27	23
Hypotonia	18	13
Tremor	15	10
Dysarthria	12	0
Psychiatric disorders		
Agitation	27	16
Insomnia	12	7
Aggression	9	0

Adverse Reactions in Pediatric Patients 6 months to Less Than 2 Years of Age

In five open-label studies including pediatric patients 6 months to less than 2 years of age with Dravet syndrome, a total of 106 patients received DIACOMIT, with 81 patients exposed for at least 6 months, and 69 patients exposed for at least 1 year. Adverse reactions in pediatric patients with Dravet syndrome who were 6 months to less than 2 years of age were similar to those seen in patients in Study 1 and Study 2.

7 DRUG INTERACTIONS

7.1 Effect of DIACOMIT on Other Drugs

CYP1A2, CYP2B6, CYP3A4, CYP2C8, CYP2C19, P-glycoprotein (P-gp) and Breast Cancer Resistance Protein (BCRP) Substrates

In vitro data show that stiripentol is both an inhibitor and inducer of CYP1A2, CYP2B6, and CYP3A4. Because of potential drug-drug interactions, consider dose adjustment of CYP1A2 substrates (e.g., theophylline, caffeine), CYP2B6 substrates (e.g., sertraline, thiotepa), and CYP3A4 substrates (e.g., midazolam, triazolam, quinidine), as clinically appropriate, when administered concomitantly with DIACOMIT.

Because of potential inhibition of enzyme/transporter activity, consider a reduction in dosage of substrates of CYP2C8, CYP2C19 (e.g., diazepam, clopidogrel), P-gp (e.g., carbamazepine), and BCRP (e.g., methotrexate, prazosin, glyburide), if adverse reactions are experienced when administered concomitantly with DIACOMIT.

Clobazam

Co-administration of DIACOMIT (which inhibits CYP 3A4 and 2C19) with clobazam results in increased plasma concentrations of clobazam (a substrate of CYP3A4) and norclobazam, the active metabolite of clobazam (a substrate of CYP2C19) [see *Clinical Pharmacology* (12.3)]. This may increase the risk of clobazam-related adverse reactions. Consider a reduction in dosage of clobazam if adverse reactions are experienced when co-administered with DIACOMIT [see *Warnings and Precautions* (5.1)].

7.2 Effect of Other Drugs on DIACOMIT

Induction-based interactions leading to decreases in DIACOMIT concentrations are possible when co-administered with a potent CYP1A2, CYP3A4, or CYP2C19 inducer, such as rifampin, phenytoin, phenobarbital and carbamazepine, as these enzymes all metabolize stiripentol. Concomitant use of strong inducers with DIACOMIT should be avoided, or dosage adjustments should be made.

7.3 CNS Depressants and Alcohol

Concomitant use of DIACOMIT with other CNS depressants, including alcohol, may increase the risk of sedation and somnolence [see *Warnings and Precautions* (5.1)].

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Exposure Registry

There is a pregnancy exposure registry that monitors pregnancy outcomes in women exposed to AEDs, such as DIACOMIT, during pregnancy. Physicians are advised to recommend that pregnant patients taking DIACOMIT enroll in the North American Antiepileptic Drug (NAAED) Pregnancy Registry. This can be done by calling the toll free number 1-888-233-2334, and must be done by patients themselves or their caregiver. Information on the registry can also be found at the website <http://www.aedpregnancyregistry.org/>.

Risk Summary

There are no adequate data on the developmental risks associated with the use of DIACOMIT in pregnant women. Administration of stiripentol to pregnant animals produced evidence of

developmental toxicity, including increased incidences of fetal malformations, increased embryofetal and pup mortality, and decreased embryofetal and pup growth, at maternal doses lower than the recommended clinical dose [see *Animal Data*].

The background risk of major birth defects and miscarriage in Dravet syndrome is unknown. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

Data

Animal Data

Oral administration of stiripentol (0, 50, 200, or 800 mg/kg/day) to pregnant mice throughout the period of organogenesis resulted in increased embryofetal mortality and decreased fetal body weights at all doses and an increased incidence of malformations at the high dose, with no evidence of maternal toxicity. The lowest effect dose for developmental toxicity in mice (50 mg/kg/day) was less than the recommended human dose (RHD) of 50 mg/kg/day on a body surface area (mg/m²) basis.

Oral administration of stiripentol (0, 50, 200, or 800 mg/kg/day) to pregnant rabbits throughout organogenesis resulted in increased embryofetal mortality at the mid and high dose and decreased fetal body weights at all doses. The mid and high doses were associated with maternal toxicity. The lowest effect dose for developmental toxicity in rabbits (50 mg/kg/day) was less than the RHD on a mg/m² basis.

Oral administration of stiripentol (0, 50, 200, or 800 mg/kg/day) to rats throughout pregnancy and lactation resulted in decreased pup survival, decreased pup body weights at birth and throughout lactation, and deficits in pup reflex development at the high dose, which was also associated with maternal toxicity. The no-effect dose for pre- and postnatal developmental toxicity in rats (200 mg/kg) was less than the RHD on a mg/m² basis.

8.2 Lactation

Risk Summary

There are no data on the presence of stiripentol in human milk, the effects on the breastfed infant, or the effects on milk production.

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for DIACOMIT and any potential adverse effects on the breastfed infant from DIACOMIT or from the underlying maternal condition.

8.4 Pediatric Use

The safety and effectiveness of DIACOMIT have been established for the treatment of seizures associated with Dravet syndrome in patients taking clobazam who are 6 months and older and weighing 7 kg or more. Use of DIACOMIT in this pediatric population is supported by 2 multicenter placebo-controlled, double-blind randomized studies in patients 3 to 18 years of age with additional pharmacokinetic and safety data in patients 6 months to less than 3 years of age [see *Clinical Studies* (14)].

The safety and effectiveness of DIACOMIT have not been established in pediatric patients below the age of 6 months or who weigh less than 7 kg.

8.5 Geriatric Use

Clinical studies of DIACOMIT in Dravet syndrome did not include patients ≥ 65 years of age to determine whether they respond differently from younger patients. The possibility of age-associated hepatic and renal function abnormalities should be considered when using DIACOMIT in patients ≥ 65 years of age [see *Clinical Pharmacology* (12.3)].

8.6 Renal Impairment

There is no formal study of the pharmacokinetics and metabolism of DIACOMIT in patients with renal impairment. However, since DIACOMIT metabolites are eliminated mainly through the kidney, administration to patients with moderate or severe renal impairment is not recommended.

8.7 Hepatic Impairment

There has been no formal study of the pharmacokinetics of DIACOMIT in patients with liver impairment. However, since the drug is mainly metabolized by the liver, administration to patients with moderate or severe liver impairment is not recommended.


10 OVERDOSAGE

There are no data concerning overdose in humans. In mice treated with high doses of stiripentol (600 to 1800 mg/kg i.p.), decreased motor activity and decreased respiration were observed. Treatment of an overdose should be supportive (symptomatic measures in intensive care units).

For management of a suspected drug overdose, contact your regional Poison Control Center.

11 DESCRIPTION

Table 4. Description

Proprietary Name	DIACOMIT
Established Name	Stiripentol
Route of Administration	Oral
Chemical Name	4,4-dimethyl-1-[3,4-(methylenedioxyphenyl)-1-pentene-3-ol]
Structural Formula	 <p>* : identifies an asymmetric carbon.</p>

Stiripentol is a white to pale yellow crystalline powder with a bitter taste; it is practically insoluble in water (at 25°C), sparingly soluble in chloroform, and soluble in acetone, ethanol, ether, acetonitrile, and dichloromethane. The melting point is approximately 75°C. The pKa is

14.2, and measurement of the partition coefficient (water-octanol) provides a Log P value of 2.94. The molecular formula is C₁₄H₁₈O₃ and the molecular weight is 234.3.

Capsules

DIACOMIT capsules contain 250 mg (size 2: pink) or 500 mg (size 0: white) of stiripentol. Capsules also contain the following inactive ingredients: erythrosine (250 mg capsule only), gelatin, indigotine (250 mg capsule only), magnesium stearate, povidone, sodium starch glycolate, titanium dioxide.

For Oral Suspension

DIACOMIT for oral suspension packets contain 250 mg or 500 mg of stiripentol. DIACOMIT packets also contain the following inactive ingredients: aspartame, carmellose sodium, erythrosine, glucose, hydroxyethylcellulose, povidone, sodium starch glycolate, sorbitol, titanium dioxide, fruit flavor (acacia, Bergamot oil, hypromellose, maltodextrin, sorbitol, and vanillin).

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

The mechanism by which DIACOMIT exerts its anticonvulsant effect in humans is unknown. Possible mechanisms of action include direct effects mediated through the gamma-aminobutyric acid (GABA)_A receptor and indirect effects involving inhibition of cytochrome P450 activity with resulting increase in blood levels of clobazam and its active metabolite.

12.2 Pharmacodynamics

There are no relevant data on the pharmacodynamic effects of DIACOMIT.

12.3 Pharmacokinetics

The following pharmacokinetic properties of stiripentol have been found in studies in adult healthy volunteers and adult patients. Systemic exposure of stiripentol increases in a greater than dose proportional manner from 500 mg to 2000 mg.

Absorption: The median time to stiripentol peak plasma concentration is 2 to 3 hours.

Distribution: Protein binding of stiripentol is 99%.

Elimination: The elimination half-life of stiripentol ranges from 4.5 to 13 hours, increasing with doses of 500 mg, 1000 mg and 2000 mg.

Metabolism: On the basis of *in vitro* studies, the main liver cytochrome P450 (CYP) isoenzymes involved in metabolism are considered to be CYP1A2, CYP2C19, and CYP3A4.

Specific Populations

The effect of age (≥ 65 years), race, renal and hepatic impairment on stiripentol pharmacokinetics is unknown [see *Use in Specific Populations* (8.5, 8.6, 8.7)]. Sex does not have a clinically significant effect on the pharmacokinetics of DIACOMIT.

Pediatric Patients: In a study of children (median age 7.3 years) with Dravet syndrome treated with DIACOMIT, valproate, and clobazam, the apparent clearance and volume of distribution of stiripentol were related to body weight. Elimination half-life increased from 8.5 hr (for 10 kg) to 23.5 hr (for 60 kg).

Drug Interaction Studies

In Vitro Studies

The metabolic pathway for stiripentol has not been clearly elucidated. Stiripentol is a substrate of several CYP enzymes, including CYP1A2, CYP2C19, and CYP3A4. Stiripentol inhibits and induces CYP1A2, CYP2B6, and CYP3A4. Stiripentol also inhibits CYP2C8, CYP2C19, and drug transporters, including P-gp and BCRP, at clinically relevant concentrations [see *Drug Interactions (7.1)*].

Clinical Studies

Antiepileptic drugs: Co-administration of clobazam with stiripentol increased concentrations of clobazam by approximately 2-fold and norclobazam (clobazam active metabolite) by 5-fold [see *Drug Interactions (7.1)*].

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

In mice, oral administration of stiripentol (0, 60, 200, or 600 mg/kg/day) for 78 weeks increased the incidences of liver tumors (hepatocellular adenoma and carcinoma) at the mid and high dose. The dose not associated with an increase in liver tumors (60 mg/kg/day) is less than the recommended human dose (RHD) of 50 mg/kg/day, based on body surface area (mg/m^2). In rats, oral administration of stiripentol at doses of up to 800 mg/kg/day (approximately 2.5 times the RHD on a mg/m^2 basis) for 102 weeks did not result in an increase in tumors.

Mutagenesis

Stiripentol was negative for genotoxicity in in vitro (Ames, HPRT gene mutation in V79 Chinese hamster cells, and chromosomal aberration in human lymphocytes) and in vivo (mouse bone marrow micronucleus) assays. Stiripentol was clastogenic in CHO cells in vitro, but only at cytotoxic concentrations.

Impairment of Fertility

Oral administration of stiripentol (0, 50, 200, or 800 mg/kg/day) to male and female rats prior to and throughout mating and continuing in females throughout organogenesis produced no adverse effects on fertility. The highest dose tested is approximately 2.5 times the RHD on mg/m^2 basis.

14 CLINICAL STUDIES

The effectiveness of DIACOMIT for the treatment of seizures associated with Dravet syndrome was established in 2 multicenter placebo-controlled double-blind randomized studies (Study 1 and Study 2), conducted according to similar protocols. To be enrolled in either study, patients were required to be 3 years to less than 18 years of age, to have Dravet syndrome (ILAE

classification of epilepsy, 1989), and to be inadequately controlled on clobazam and valproate, with at least 4 generalized clonic or tonic-clonic seizures per month despite optimized therapy.

Eligible patients were enrolled in a 1-month baseline period during which they continued to receive their optimized antiepileptic treatment. Following this 1-month baseline, patients were randomly allocated to receive either DIACOMIT (fixed dose of 50 mg/kg/day in divided doses with no dose titration) or placebo, added to their treatment with clobazam and valproate. Duration of double-blind treatment was 2 months. The frequency of generalized clonic or tonic-clonic seizures during the study was recorded by patients and/or their caregivers, using a diary. Although patients with Dravet syndrome have several different types of seizures, only generalized clonic or tonic-clonic seizures were recorded, as other seizure types can be difficult to recognize by patients and/or their caregivers as seizures.

The primary efficacy endpoint for both studies was the responder rate. A responder was defined as a patient who experienced a greater than 50% decrease in the frequency (per 30 days) of generalized clonic or tonic-clonic seizures during the double-blind treatment period compared to the 4-week baseline period (i.e., placebo run-in). The mean change from baseline in frequency of generalized clonic or tonic clonic seizures was also evaluated.

In Study 1 (n=41), 21 patients were randomized to DIACOMIT, and 20 patients to placebo. In Study 2 (n=23), 12 patients were randomized to DIACOMIT, and 11 patients to placebo. In both studies, the demographic and baseline clinical characteristics were similar between the treatment groups.

Table 5 summarizes the results of the primary endpoint for DIACOMIT in each study.

Table 5. Efficacy Results in the Intent-to-Treat Population in Study 1 and Study 2

	Study 1 N=41		Study 2 N=23	
	DIACOMIT N=21	Placebo N=20	DIACOMIT N=12	Placebo N=11
Responder Analysis ^a				
No of responders/total (Responder Rate) [95% CI]	15/21 (71%) [52% – 91%]	1/20 (5%) [0.0% – 15%]	8/12 (67%) [40% – 93%]	1/11 (9.1%) [0.0%– 26%]
p-value ^b	<0.0001		0.0094 ^e	
Percentage Change from Baseline in Seizure Frequency ^c				
n	20	16	11	9
Mean ± SD	-69% ± 42%	7.6% ± 38%	-74% ± 27%	-13% ± 62%
Median	-91%	7.4%	-81%	-27%
Min – Max	-100% – 28%	-75% – 65%	-100% – -33%	-87% – 140%
p-value ^d	0.0002		0.0056 ^e	

^a Responder is defined as a patient with a greater than 50% decrease in frequency of generalized tonic-clonic or clonic seizures

^b Fisher Exact Test

^c Frequency of generalized tonic-clonic or clonic seizures during month 2

^d Wilcoxon Test with two-sided t-approximation


^e Nominal p value, as Study 2 was stopped early

CI=confidence interval; SD=standard deviation.

In both studies, the responder rate (primary efficacy endpoint) was significantly greater for DIACOMIT than for placebo. DIACOMIT was also superior to placebo for the reduction in mean frequency of generalized clonic or tonic-clonic seizures. In Study 1 and Study 2, respectively 43% and 25% of patients reported no generalized clonic or tonic-clonic seizure for the duration of the study.

Figure 1 displays the percentage of patients by category of percent reduction in tonic-clonic and clonic seizure frequency during month 2 of the treatment period compared to baseline (per 30 days) in Study 1 and Study 2 (pooled).

Figure 1.: Proportion of Patients by Category of Seizure Response for DIACOMIT and Placebo in Study 1 and Study 2 Pooled, Baseline to 2nd Month of Treatment (per 30 days).



The effectiveness of DIACOMIT for the treatment of seizures associated with Dravet syndrome in patients 6 months of age to less than 3 years of age was extrapolated from the demonstration of effectiveness in patients 3 years to less than 18 years of age in Study 1 and Study 2.

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

DIACOMIT Capsules

- 250 mg: size 2, pink, and imprinted with “Diacomit” and “250mg” are supplied as follows:
Bottles of 60 NDC 68418-7939-6
- 500 mg: size 0, white, and imprinted with “Diacomit” and “500mg” are supplied as follows:
Bottles of 60 NDC 68418-7940-6

For Oral Suspension

- 250 mg: pale pink fruit flavored powder packaged in packets are supplied as follows:
Cartons of 60 NDC 68418-7941-6
- 500 mg: pale pink fruit flavored powder packaged in packets are supplied as follows:
Cartons of 60 NDC 68418-7942-6

16.2 Storage and Handling

Store in a dry place at 20°C to 25°C (68°F to 77°F); excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature]. Store in original package to protect from light.

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Medication Guide and Instructions for Use).

DIACOMIT Oral Capsule Administration

Inform patients or caregivers that DIACOMIT capsules must be swallowed whole with a glass of water during a meal. Capsules should not be broken or opened.

DIACOMIT For Oral Suspension Administration

DIACOMIT should be mixed in a glass of water and should be taken immediately after mixing during a meal [*see Instructions for Use*].

Somnolence

Advise patient or caregivers that somnolence may occur, and may require a decrease in the dose of clobazam [*see Warnings and Precautions (5.1)*]. Also, advise the patients and their caregivers to avoid alcohol consumption during DIACOMIT treatment [*see Drug Interactions (7.3)*].

If applicable, caution patients about hazardous machinery, including automobiles, until they know how DIACOMIT affects them.

Decreased Appetite and Decreased Weight

Advise patients or caregivers that decreased appetite is frequent and nausea and vomiting can also occur during DIACOMIT treatment, which can cause loss of weight [*see Warnings and Precautions (5.2)*].

Withdrawal Symptoms

Advise patients or caregivers that abrupt withdrawal of DIACOMIT may increase their risk of seizures or status epilepticus [*see Dosage and Administration (2.5) and Warnings and Precautions (5.4)*]. Instruct patients or caregivers to not discontinue use of DIACOMIT without consulting with their healthcare provider.

Neutropenia and Thrombocytopenia

Advise patients or caregivers of the risk of neutropenia and thrombocytopenia and the importance of hematologic testing, which should be obtained prior to starting treatment with DIACOMIT and then every 6 months. *[see Warnings and Precautions (5.3)]*.

Suicidal Thinking and Behavior

Counsel patients, their caregivers, and their families that AEDs, including DIACOMIT, may increase the risk of suicidal thoughts and behavior and advise them of the need to be alert for the emergence of worsening of symptoms of depression, any unusual changes in mood or behavior, or the emergence of suicidal thoughts, behavior, or thought of self-harm. Patients or caregivers should report behaviors of concern immediately to healthcare providers *[see Warnings and Precautions (5.5)]*.

Use in Pregnancy

Advise patients to notify their healthcare provider if they become pregnant or intend to become pregnant during DIACOMIT therapy. Encourage patients to enroll in the NAAED Pregnancy registry if they become pregnant. This registry is collecting information about the safety of AEDs during pregnancy *[see Use in Specific Populations (8.1)]*.

Use in Nursing

Instruct patients to notify their physician if they are breast feeding or intend to breast feed during therapy *[see Use in Specific Populations (8.2)]*.

DIACOMIT Capsules and DIACOMIT for Oral Suspension manufactured by:
BIOCODEX
1, avenue Blaise Pascal
60000 BEAUVAIS
France